

Datasheet

Anti-Myostatin Clone Myo 2/1A

Product Name	Anti Human Myostatin Myo 2/1A
Catalogue Number	Myo2/1A
Clone, Isotype	Myo 2/1A
Format	IgG
Tested Applications	WB, ICC

Description:

Myostatin is a well-characterized negative regulator of skeletal muscle and can inhibit myogenesis and stimulate adipogenesis. Clone Myo 2/1A has been shown to have the reverse effect, up-regulate myogenesis and down-regulate adipogenesis. (Artaza, N et al.)

Product Details:

Form in stock: IgG, purified – 1.0 mg/mL. Also available as unpurified supernatant.

Host: Mouse

Specificity: Recognizes the 113 amino acid carboxy-terminal fragment of Myostatin protein.

Human Histology positive control: Muscle fibre



Storage: Store at +4°C or -20°C. Avoid repeated freezing and thawing.

Shelf life: 18 months from date of dispatch.

Regulatory/ Restrictions: For research and commercial purposes.

Applications	Suggested Dilution
Western Blot	1:500 ¹
Immunocytochemistry	100 µg/mL ¹


Applications:



Clone Myo 2/1A used to detect Myostatin expression by ICC

Image caption: Effect of recombinant Mst-113 protein and anti-Mst antibody on myogenesis in differentiating C3H 10T(1/2), assessed by using myogenin immunocytochemical staining. Cells were incubated with medium alone (C, control), graded concentrations of recombinant Mst-113 or anti-Mst antibody (Mst Ab) for 1 wk. A, Immunocytochemical staining using an antibody against myogenin; quantitative image analysis of stained cells is shown in B. *Asterisks* denote the *P* values for the statistical comparison of treatment group against medium control. No 1st Ab, Cells not treated with first antibody. Magnification, ×400 (Artaza, N et al.)


Dilution used: 100 µg/mL



Clone Myo 2/1A used to detect Myostatin expression by Western Blot with detection at 52 kDa

Image caption: A: Luminol detection of Western blots with monoclonal antibody against Mst at 3 d after transfection (Artaza, N et al.)

Dilution used: 1:500



Clone Myo 2/1A used to detect C/EBP α to assess adipogenic differentiation C3H 10T(1/2) by IHC

Image caption: B, Cells were incubated for 4 d (C/EBP α) and 1 wk (adiponectin) with graded concentrations of recombinant Mst-113 or anti-Mst antibody, and plates were divided for either trypsination and Western blot on the cell extracts (30 μ g/lane) with an antibody against C/EBP- α (Artaza, N et al.)

Dilution used: 100 μ g/mL

References:

1. Artaza, J.N., Bhasin, S., Magee, T.R., Reisz-Porszasz, S., Shen, R., Groome, N.P., Fareez, M.M., Gonzalez-Cadavid, N.F. (2005) Myostatin Inhibits Myogenesis and Promotes Adipogenesis in C3H 10T(1/2) Mesenchymal Multipotent Cells. *Endocrinology*; 146 (8): 3547-3557.