Datasheet

Anti-Spike (RBD) Antibody_BSV-COV-AB-44

BioServUK offers a series of different monoclonal antibodies for the detection of the Spike protein from SARS-CoV-2. All antibodies come from hybridoma cell lines that were generated by immunization with the recombinant full-length Spike protein (S) or the receptor-binding domain (RBD).

This antibody has been validated by ELISA and is specifically directed against an epitope that is located on the RBD. Additionally, ELISAs with directly coated antigens proved that this antibody also binds to Spike protein B.1.1.7 and different RBD variants (B.1.1.7, B.1.1.28.1, B.1.1.7 E484K and B.1.351).

All antibodies are produced exclusively under serum-free conditions from hybridoma and purified through one-step purification with Protein-A affinity chromatography.

Product Code:	BSV-COV-AB-44
Host:	Mouse
Clonality:	Monoclonal
Isotype:	IgG
Subclass:	mIgG2ak
Formulation:	Liquid, PBS, pH 7.4, 0.2 µm sterile filtered
Concentration:	≥ 0.5 mg/ml
Purity:	≥ 90% (via analytical CGE under reducing conditions)
Conjugate:	Unconjugated
EC_{50} RBD*:	2.117 nM
EC_{50} S1*:	1.117 nM
EC_{50} S*:	0.424 nM
Specificity:	S (tested for: Wuhan and B.1.1.7) RBD (tested for: Wuhan, B.1.1.7, B.1.1.28.1, B.1.1.7 E484K and B.1.351)

*EC50 values for 20 nM of coated antigen (Wuhan origin)

The product is for research use or for further manufacturing only.

For UK, Europe and ROW orders:
Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way, Rotherham
S60 5XA
+44 114 224 2257
info@bioservuk.com

For Americas Orders:
Anatrace
434 West Dussel Drive
Maumee, OH 43537
+1 800 252 1280
customerservice@anatrace.com

Purity (analytical CGE, under reducing conditions):



Peak	Size [kDa]	% of Total	Peak Identification
1	2.9	0.0	Unknown
2	4.5	0.0	Lower Marker
3	5.6	0.0	System Peak
4	7.9	0.0	System Peak
5	19.3	0.3	Unknown
6	25.3	Peak Value 1	Light Chain
7	58.2	Peak Value 2	Heavy Chain
8	67.2	0.5	Unknown
9	240.0	0.0	Upper Marker

Summation of peak values 1 and 2 results in a purity of ≥ 90%

For UK, Europe and ROW orders:
Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way, Rotherham
S60 5XA
+44 114 224 2257
info@bioservuk.com

For Americas Orders:
Anatrace
434 West Dussel Drive
Maumee, OH 43537
+1 800 252 1280
customerservice@anatrace.com

Protein Activity (ELISA):



	154 nM RBD	77 nM RBD	39 nM RBD	19 nM RBD
EC_{50} [nM]	0.5171	0.6327	0.8365	2.516

For ELISA analyses immunoassay plates were coated with different concentrations of SARS-CoV-2 RBD (19, 39, 77 and 154 nM). Antibody BSV-COV-AB-44 was applied at concentrations ranging from 26.67 nM to 0.026 nM.

Bound antibody BSV-COV-AB-44 was detected using Goat Anti-Mouse-IgG-HRP (10 ng/ml) and TMB. Determined EC_{50} of antibody BSV-COV-AB-44 for 20 nM RBD is 2.117 nM.



	84 nM S1	42 nM S1	21 nM S1	10 nM S1
EC_{50} [nM]	0.4551	0.5326	0.9286	2.430

For ELISA analyses immunoassay plates were coated with different concentrations of SARS-CoV-2 S1 (10, 21, 42 and 84 nM). Antibody BSV-COV-AB-44 was applied at concentrations ranging from 26.67 nM to 0.026 nM.

Bound antibody BSV-COV-AB-44 was detected using Goat Anti-Mouse-IgG-HRP (10 ng/ml) and TMB. Determined EC_{50} of antibody BSV-COV-AB-44 for 20 nM S1 is 1.117 nM.

For UK, Europe and ROW orders:
Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way, Rotherham
S60 5XA
+44 114 224 2257
info@bioservuk.com

For Americas Orders:
Anatrace
434 West Dussel Drive
Maumee, OH 43537
+1 800 252 1280
customerservice@anatrace.com



	29 nM S	14 nM S	7 nM S	3.6 nM S
EC_{50} [nM]	0.5057	0.5967	0.7486	1.715

For ELISA analyses immunoassay plates were coated with different concentrations of SARS-CoV-2 Spike spike protein (3.6, 7, 14 and 29 nM). Antibody BSV-COV-AB-44 was applied at concentrations ranging from 26.67 nM to 0.026 nM.

Bound antibody BSV-COV-AB-44 was detected using Goat Anti-Mouse-IgG-HRP (10 ng/ml) and TMB. Determined EC_{50} of antibody BSV-COV-AB-44 for 20 nM Spike protein is 0.424 nM.

Binding activity to SARS-CoV-2 Spike protein and receptor-binding domain variants (ELISA):



Binding activity of BSV-COV-AB-44 to coated SARS-CoV-2 Spike protein (Wuhan origin vs B.1.1.7) in comparison to 19 other monoclonal mouse hybridoma antibodies targeting the Spike protein.

Bound antibodies were detected using anti-mouse IgG antibodies conjugated to HRP in an in-house SARS-CoV-2 IgG ELISA.

For UK, Europe and ROW orders:
Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way, Rotherham
S60 5XA
+44 114 224 2257
info@bioservuk.com

For Americas Orders:
Anatrace
434 West Dussel Drive
Maumee, OH 43537
+1 800 252 1280
customerservice@anatrace.com



Binding activity of BSV-COV-AB-44 to different coated SARS-CoV-2 receptor binding domain variants (RBD Wuhan, B.1.1.7, B.1.1.28.1, B.1.1.7, E484K and B.1.351) in comparison to 12 other monoclonal mouse hybridoma antibodies targeting the RBD and a negative control antibody (NEG_CON).

Bound antibodies were detected using anti-mouse IgG antibodies conjugated to HRP in an in-house SARS-CoV-2 IgG ELISA. The negative control is a monoclonal mouse hybridoma antibody which does not bind the RBD of SARS-CoV-2.

For UK, Europe and ROW orders:
Calibre Scientific UK
Unit 5A, R-Evolution @
The Advanced Manufacturing Park
Selden Way, Rotherham
S60 5XA
+44 114 224 2257
info@bioservuk.com

For Americas Orders:
Anatrace
434 West Dussel Drive
Maumee, OH 43537
+1 800 252 1280
customerservice@anatrace.com